

LOUISIANA RECOMMENDATIONS FOR CONTROL OF INSECTS ON MAYHAW

Insect	Material	Rate	Restrictions
Aphids (except Woolly apple aphid)	Provado 1.6 F	2.0-4.0 fl ozs/100 gals or 4.0-8.0 fl ozs/acre	Provado: PHI=7 days; apply post-bloom; highly toxic to bees.
	or Mustang Max	0.21-0.32 fl oz/50 gals or 1.28 fl ozs/acre	Mustang Max: PHI=14 days; allow 7 days between applications.
Leafminers	SpinTor 2SC	1.0-2.5 fl ozs/100 gals or 4.0-10.0 fl ozs/acre	SpinTor: PHI=7 days; do not apply more than 29 fl ozs/acre/year.
Leafhoppers	Provado 1.6F	1.0-2.0 fl ozs/100 gals or 4.0-8.0 fl ozs/acre	Provado: PHI=7 days; apply post-bloom; highly toxic to bees.
	or Mustang Max	0.21-0.32 fl oz/50 gals or 1.28 fl ozs/acre	Mustang Max: PHI=14 days; allow 7 days between applications.
Aphids Plum curculio	Actara	4.5 ozs/acre	Actara: PHI=35 days; highly toxic to bees; do not make more than 1 pre-bloom application.
	or Mustang Max	0.21-0.32 fl oz/50 gals or 1.28 fl ozs/acre	Mustang Max: PHI=14 days; allow 7 days between applications.
Spider mites	Savey 50DF	3.0-6.0 ozs/acre	Savey: PHI=28 days; apply at the beginning of mite infestations. Savey is not effective against adult mites. Limit to 1 application/year. Do not use in home plantings.

Warning: Re-entry time for worker entering treated areas should be strictly observed. Read the label for this information.