


# ***Louisiana*** **Home Citrus** *Production*



**LSU**  
**AgCenter**  
Research & Extension



## Site Selection

A well-drained soil, high in organic matter and slightly acid to neutral, is desirable for citrus. The site should provide full sunlight. In most cases, homeowners are limited to sites in the yard where citrus can be planted. The site with the most sun should be selected for a citrus planting. Citrus trees require good drainage. Trees planted on heavy clay soils with poor internal drainage should be planted on a mound or row 8 to 12 inches high.

## Planting

The best time to plant citrus trees is in January or February. Trees planted after December can withstand a freeze in February better than earlier-planted trees. A 2- to 4-foot tree with three to four well-developed upward-growing side branches 18 to 24 inches above the ground is the ideal tree for home plantings.

The root system of the young trees should be inspected before planting. Trees that have bent or circled root systems will be stunted and grow poorly. Nurserymen and growers often called this condition 'j' rooted. Prune any damaged roots back beyond the damaged area. If container-grown trees have a tap root curled in the bottom of the container, cut this root off at the point where it begins to curl. The tree will grow a new tap root. Separate and trim the roots of container trees that are root bound.

Dig a hole slightly larger than the container. Place the tree in the hole at the same depth it was growing previously. If holes are dug too deeply trees may settle after watering. Trees set too deeply may die. Container trees should have the top of the soil flush with the top of the hole. Bare-rooted trees should have soil placed underneath them in a manner to allow the spreading of the roots in a natural position with no bending or crimping.

Before completion of backfilling, add water to settle the soil and eliminate air pockets around the roots. After watering, fill the hole to completion, and water again. Construct a ridge around the complete circumference of the tree to hold water during subsequent watering.

## Spacing

Different types of citrus require different spacing. In most cases, homeowners are limited to the space in their yards for citrus trees. The site that will provide the most space should be selected. Trees planted too closely together or against buildings will have limited sunlight and air movement that will

restrict the growth and yield of the tree and enhance the development of pests. Spraying for pests, pruning and harvesting the fruit are difficult on trees planted too closely together.

Navel oranges, grapefruit and other oranges are the most vigorous citrus trees. They require at least a 30- to 40-foot diameter circle. (Example: Allow at least 15 to 20 feet from any building or large tree on each side of the navel orange, grapefruit or other round oranges). A satsuma is not as vigorous as oranges or grapefruit and requires a 20- to 30-foot circle in diameter; kumquats and lemons need only a 15- to 20-foot diameter circle.

## Rootstocks

The best citrus rootstock for the Louisiana home citrus orchard is trifoliata (*Poncirus trifoliata* Rubidoux). It is the most cold hardy of the citrus rootstock, resistant to rots and tolerant of wet soils. Homeowners should ask the nursery to obtain trees on trifoliata, but they are at the mercy of the nursery on the rootstocks of the citrus trees they buy.

One of the major rootstocks used in the Louisiana citrus nursery trade is Swingle citrumello. This very vigorous rootstock produces a large budded tree in one year. Trees budded on Swingle rootstocks are vigorous and produce good crops of high quality citrus. The main drawback of using Swingle rootstock is that it is not as cold hardy as trifoliata.

In the last several years, several nurseries have been grafting citrus trees on a dwarf rootstock known as Flying Dragon Trifoliata. Dwarf trees are ideal for homeowners with limited space. Good crops of oranges and satsuma can be made in a circle only 10 feet in diameter. Ask your local garden centers and nurseries to obtain trees budded on the dwarf Flying Dragon Trifoliata rootstocks.

## Fertilizer

Citrus trees require annual fertilization for good growth and high yields of good size, high quality fruit. Newly set trees should not be fertilized until they show signs of growth, usually six weeks after they are set in the spring (mid March). On newly planted trees, apply 1/2 pound of 8-8-8 or 13-13-13 per tree in mid March.

After the second year, fertilize citrus trees in late January or early February. Apply 1 to 1 1/2 pounds of 8-8-8 or 13-13-13 per year of tree age up to 12 years. Increase the rate of fertilizer 1 -1 1/2 pound per year as the tree gets older. A one-year-old tree gets 1 to 1 1/2 pounds of 8-8-8 or 13-13-13, and a 5-year-old tree gets from 5 to 7 1/2 pounds. The fer-

tilizer rate is increased each year until the tree is 12 years old. Trees 12 years old and older are at the top limit of fertilizer and should receive 12 to 18 pounds of 8-8-8 or 13-13-13 per tree.

A simple fact to remember is that one pint of 13-13-13 weighs about 1 pound and a quart weighs 2 pounds. (Example: A one-year-old tree will require 1 pound or one pint of 13-13-13, and a 4-year-old tree will require 4 to 6 pounds or two to three quarts of 13-13-13.) Table 1 shows the fertility schedule for citrus trees.

A small amount of nitrogen fertilizer (1/4 lb of Ammonium Nitrate or Ammonium Sulfate/year/age of tree or 1/2 lb of Calcium Nitrate/year/age of tree) should be applied to bearing trees (trees 4 years and older) in late May or June. The rate of fertilizer is increased each year until the tree is 12 years old. This helps the tree make adequate vegetative growth and size the fruit. The additional nitrogen will also encourage the tree to set a crop the following year. Nitrogen fertilizer should not be applied after the end of June. Fertilizer applied after the end of June will

decrease cold hardiness and delay fruit from ripening. Table 2 shows the summer nitrogen fertilizer schedule for citrus trees.

Broadcast the fertilizer beyond the spread of the limbs where most of the feeder roots occur. A general rule when fertilizing trees is to put your left shoulder near the outer branches and hold the can of fertilizer in your right hand. You can simply walk around the tree and evenly spread the fertilizer in a 12- to 18-inch band on the outer branches of the tree. This technique will ensure that the fertilizer is placed a safe distance from the tree.

Avoid fertilizing citrus trees after the end of June. Late fertilization will encourage vigorous growth, delay fruit maturity and decrease the cold hardiness of the tree. Homeowners are also encouraged to pay strict attention to the amounts of fertilizer applied to citrus trees. Amounts above those recommended will encourage vigorous growth, delay fruit maturity and decrease cold hardiness. Vigorous growth can result in extensive freeze damage or death of the trees, even in a moderate freeze.

Tree Age	Time of Year	Amount of Fertilizer per Tree
Year of transplanting	Mid March 6 weeks after transplanting	1/2 lb. of 8-8-8 or 13-13-13
First year	Late January-early February	1-1 1/2 lb. of 8-8-8 or 13-13-13
Second year	Late January-early February	2-3 lbs. of 8-8-8 or 13-13-13
Third year	Late January-early February	3-4 1/2 lbs. of 8-8-8 or 13-13-13
Fourth year	Late January-early February	4-6 lbs. of 8-8-8 or 13-13-13
Fifth year	Late January-early February	5-7 1/2 lbs. of 8-8-8 or 13-13-13
Sixth year	Late January-early February	6-9 lbs. 8-8-8 or 13-13-13
Increase the rate of fertilizer 1 - 1 1/2 lbs. of 8-8-8 or 13-13-13 per year as the tree gets older.		
Twelfth year and older	Late January-early February	12-18 lbs. of 8-8-8 or 13-13-13

Tree Age	Time of Year	Amount of Fertilizer per Tree
Fourth year Bearing age	Late May or June	1 lb AmNO <sub>3</sub> or AmSO <sub>4</sub> or 2 lb CaNO <sub>3</sub>
Fifth Year	Late May or June	1 1/4 lb AmNO <sub>3</sub> or AmSO <sub>4</sub> or 2 1/2 lb CaNO <sub>3</sub>
Sixth Year	Late May or June	1 1/2 lb AmNO <sub>3</sub> or AmSO <sub>4</sub> or 3 lb CaNO <sub>3</sub>
Seventh Year	Late May or June	1 3/4 lb AmNO <sub>3</sub> or AmSO <sub>4</sub> or 3 1/2 lb CaNO <sub>3</sub>
Increase the rate of AmNO <sub>3</sub> or AmSO <sub>4</sub> 1/4 lb (CaNO <sub>3</sub> 1/2 lb) per year age of tree as the tree gets older		
Twelfth Year and Older	Late May or June	3 lbs AmNO <sub>3</sub> or AmSO <sub>4</sub> or 6 lb CaNO <sub>3</sub>

## Pruning Trees

Good nursery trees usually have a framework already developed when purchased.

Homeowners should try to select a 2- to 4-foot tree. Trees should be pruned after planting and before growth starts in the spring. The top of the tree should be removed 18 to 24 inches from the ground at a site where there are three or four evenly spaced wide-angle lateral branches with an upward-growing pattern that has developed. All growth developing below this framework should be removed.


The primary purpose of pruning young non-bearing trees is to shape the tree so that scaffold branches will be well distributed. This initial pruning helps bring the tops of the plants and the root system into balance. It also stimulates good scaffold branch development. If scaffold branches are selected properly, the weak narrow crotches and downward growing branches can be eliminated and future breakage under heavy fruit loads can be avoided. Proper selection of scaffold branches also will reduce the large pruning cuts in the future years.

After the selection of the scaffold branches, only limited pruning is needed on citrus. All pruning on older trees should be done in January and February. Pruning trees of bearing age is practiced to thin out thick growth to spray and harvest easily. The removal of long vigorous growing shoots that stick up at the top of the tree will help to control the size of the tree. These long shoots should be traced to where they originate on larger branches and cut off flush at the point of attachment. Dead branches, branches crossing over each other, water sprouts arising from the center of the tree and branches touching the ground should be removed. When pruning, cut all limbs flush at the point of attachment.

Freeze-damaged trees should not be pruned until the extent of cold damage has been determined. Normally, the damage is not evident until July and August after the second flush of growth. Pruning a freeze-damaged tree consists of removing the dead wood to the point where the live wood starts.

## Varieties of Citrus

Citrus varieties vary with cold hardiness, and are listed below from hardy to tender.



## Recommended Varieties of Citrus

### Satsuma

The satsuma is the traditional type of citrus grown by Louisiana homeowners. The fruit is easy to peel, has few seeds and separates easily into segments. The fruit turns from green to yellow as it ripens and to orange at full maturity. The fruit is edible when it shows some yellow color. This allows homeowners to harvest and eat the fruit for a long period.

### Owari Satsuma

Owari is the most widely grown satsuma. The fruit is medium to small, seedless, has excellent quality and matures from early to mid November, but can be harvested through early December. The trees are vigorous and have a willowy growth habit. An Owari satsuma is recommended for home orchards in Zones I, II and the southern part of Zone III.

### Armstrong Early Satsuma

Fruit reaches maturity in late September and October. The fruit is large but has only fair quality. It becomes puffy quickly and will split badly during heavy rainfall. Armstrong satsumas should be harvested by early November for best quality. The trees are dwarfed and have an upright growth habit. Armstrong Early is recommended for home orchards in Zones I and II.

## Brown's Select Satsuma

This variety produces medium to large fruit. The fruit matures mid October to early November, several weeks ahead of Owari Satsuma. The fruit keeps well on the trees without becoming puffy. The trees are large and have an open spreading branching pattern. Brown's Select is recommended for home orchards in Zones I and II.

## Kimbrough Satsuma

Kimbrough was released as a cold-hardy Satsuma; unfortunately, its cold hardiness is no better than that of Owari. The trees of Kimbrough are large, strong, spreading and very productive. The fruit is generally larger than Owari and matures in early to mid November. The fruit stores well on the tree after maturity. Kimbrough may be hard to find in the nursery trade. This variety is recommended for Zones I and II.

## Louisiana Early and Early St. Ann Satsuma

Louisiana Early and Early St. Ann are two recently released satsuma varieties from the LSU AgCenter. They produce medium to large fruit that matures in early September through mid October. The overall quality and productivity are better than Early Armstrong. Fruit puffiness ranges from none to slight on both varieties. The fruit does not hold well on the tree for an extended period. The medium-sized trees have a spreading growth pattern. The availability of both of these varieties may be limited in the nursery trade. Louisiana Early and Early St. Ann are recommended for trials in home orchards in Zones I and II.

## Sweet Oranges

**Louisiana Sweet:** Louisiana Sweet is the traditional round orange grown in South Louisiana. It is a medium to large fruit with a rich flavor and lots of seeds. The fruit matures in December. After reaching maturity, the fruit splits badly and drops. It is the most cold hardy of the sweet oranges. The trees are very vigorous and have thorns. This variety is recommended for home orchards in Zone I and the southern part of Zone II.

**Washington Navels:** The most distinctive feature of this orange is the presence of a navel, a small rudimentary, secondary fruit embedded in the end of the fruit. The fruit is large, has excellent quality and matures in late November and December. Distinctive characteristics of navel oranges include deep orange color, thin skin, ease of peeling, separation of the segments, high sugars, abundance of juice and seedlessness. The fruit will drop after full maturity, so it's best to harvest navels by end of January. The juice often becomes bitter when stored and should be con-

sumed shortly after squeezing. Navel orange trees are usually less productive than other sweet oranges. The navel orange should be included in home orchards in Zone I and the southern part of Zone II.

**Hamlin Sweet:** This is the most widely grown of the early round oranges. It matures in early December, but it can hold on the tree through February. The fruit is medium to small and has few or no seeds. Trees are medium large, moderately vigorous and fairly cold tolerant. Hamlin is recommended for home orchards in Zone I.

**Pineapple Sweet:** Pineapple Sweet is a medium orange that matures in early December and has a pineapple flavor. The fruit is very seedy. The trees are vigorous good annual producers. Pineapple Sweets are recommended for home planting in Zone I.

**Plaquemines:** Plaquemines is a seedless, low acid, round orange. It is a bud sport of Pineapple found at Magnolia Orange Grove, Port Sulphur, La., and released as a variety by LSU in 1948. The fruit of Plaquemines is medium, seedless and matures in January. The trees are vigorous good annual producers. Plaquemines is recommended for home orchards in Zone I.

**Valencia:** Valencia is the most widely planted orange in the world. The fruit is medium-large with few or no seeds. It ripens in April and has excellent quality through June. Juice is abundant, and flavor is excellent. Trees are vigorous, upright and prolific. It is subject to heavy fruit drop after a freeze. This variety is recommended for home plantings only in the southernmost parts of Zone I.

**Ambersweet:** Ambersweet is a sweet orange hybrid released in 1989 by the USDA Horticultural Research Station in Orlando. It is a hybrid of Clementine tangerine by Orlando tangelo crossed with a seedling mid season sweet orange. Moderately cold tolerant, this early season orange is of peak harvesting quality from October through December. Ambersweet fruit are medium-sized, slightly pear shaped, low in acid with good juice and flesh quality. Fruit in a home planting can have up to 30 more seeds in mixed plantings. Homeowners are encouraged to ask local nursery and garden centers to obtain trees of Ambersweet. It is recommended for trial in home planting in Zone I.

## Moro Blood Oranges

This is a medium to medium-large, round, sweet orange with very few seeds. It is characterized by red coloration in the flesh and peel. It reaches maturity in late December and holds on the tree very well. The riper the fruit, the redder the fruit's pigment. The trees are of medium vigor and size with a spreading round topped shape. Homeowners are encouraged

to ask local nurseries and garden centers to obtain trees of Moro Blood oranges. Moro Blood oranges are recommended for trial in home orchards in Zone I.

## Grapefruit

**Ruby Red:** Ruby Red is the most widely planted grapefruit variety in Louisiana. It matures in December and holds well on the tree through May. The fruit is medium to large and has only a few seeds, light yellow skin color at maturity with a red blush, especially where fruit touch one another. It is recommended for home orchards in Zone I and southern part of Zone II.

**Rio Red:** Rio Red produces larger fruit than Ruby Red. The fruit is yellow at maturity with a red blush. This variety is earlier than Ruby Red and can be harvested from early November through May. It is recommended for trials in home orchards in Zone I and southern part of Zone II.

## Kumquats

**Nagami Kumquats:** The Nagami kumquat produces oblong fruit with a smooth rind, deep orange color and acid juice. It ripens from mid October to February. The fruit contains seeds. The Nagami trees are vigorous with a round, bushy top. Its growth pattern makes it adaptable to hedge and corner plantings. Nagami kumquats are cold hardy and recommended for home orchards in Zones I, II and the southern part of Zone III.

**Meiwa Kumquats:** The Meiwa kumquats produce round fruit with sweet pulp. The trees are less vigorous than Nagami. Meiwa kumquats are cold hardy and recommended for home orchards in Zones I, II and the southern part of Zone III.

## Lemons

**Myers Lemons:** Myers is the only lemon recommended for Louisiana since it does possess a small degree of cold hardiness. It ripens in mid October and holds on the tree until December. It is better when grown from a rooted cutting. It has a strong tendency to bloom and set fruit throughout the year. This makes it an excellent tree for a protected area near a window or door. It is recommended for homeowners in Zone I and in protected areas in the southern part of Zone II.

## Other Citrus

**Ponkan Mandarin:** The fruit of Ponkan mandarin is medium to large and develops a deep orange color when mature. It matures in mid December and should be harvested by mid January. It has few seeds and a honey-type mandarin flavor. If left on the tree

too long, it tends to become puffy. The tree is moderately vigorous and has a very upright growth habit and weak crotches. It has a tendency to bear alternately and suffers from limb breakage in heavy crop years. Ponkan Mandarin is recommended for home orchards in Zone I.

**Orlando Tangelo:** Orlando tangelo is a hybrid between a Duncan grapefruit and Dancy tangerine. The fruit is medium in size and has high juice content. Fruit has deep orange color in late December with good quality until mid January. Orlando tangelo is recommended for home orchards in Zone I.

**Dancy Tangerine:** The most extensively planted tangerine variety; the fruit ripens in mid December to February. Fruit is medium in size and has a slightly flattened shape. The fruit color is deep orange-red at maturity. The fruit dries out when left on the tree for a long time after becoming ripe. Dancy tends to overbear and has brittle wood, resulting in frequent limb breakage. This variety is recommended for planting only in Zone I.

**Robinson Tangerine:** Robinson tangerine is a cross between Clementine mandarin and Orlando tangelo. This variety produces medium to large fruit with deep orange red flesh of excellent quality. It peels easily. It matures in mid October. It produces best when planted with Orlando tangelo or Sunburst tangerine since it will not set fruit with its own pollen. It has brittle wood and a tendency to set fruit near the end of its limbs, resulting in limb breakage with large crops. This variety is recommended for home planting in Zone I.

**Sunburst Tangerine:** Sunburst tangerine is a cross between Robinson and Osceola tangerine. The fruit ripens in late November and December and holds well on the tree. Fruit are reddish-orange and are high quality with good flavor. It requires cross pollination with Orlando tangelo for good fruit set. It is very susceptible to rust mite injury. Sunburst tangerine is recommended for trial plantings in home orchards in Zone I.

## Picking

Citrus is the ideal fruit for the home grower because it can remain on the tree in a good state for a number of months. Satsumas can be harvested in the green-yellow stage as early as late September early October, while oranges, grapefruit and other citrus are not ready to harvest until they reach the full color stage – starting in late November through early December.

Home growers can start picking satsumas and kumquats as soon as the fruit starts turning from yellow to green. A good rule is to taste a satsuma or

kumquat; and if you like the taste, then start picking a few of the fruit each week. The fruit left on the tree will turn from yellow to orange, develop a loose skin and become sweeter. Just be aware that satsumas will not separate cleanly from the stem. They should be clipped from the stem to prevent tearing the skin of the fruit.

Citrus fruit left on the tree will develop more color and improve in quality with exposure to cold temperatures. It takes temperature in the mid- to low 20s five to 10 hours to freeze the fruit. All fruit should be picked by the end of January or early February. Fruit left on trees after this time will reduce the blooms and fruit set for next year.

## Citrus Insects

### Citrus Red Mite (*Panonychus citri*)

The citrus red mite has a small red body with several white hairs (setae) arising from the back and sides of the abdomen. Each female can lay two to three eggs a day and may lay 20 to 50 eggs. The mite eggs are red with white setae in the top center. They can develop from egg to egg in 12 days. Populations increase in spring, late summer and early fall in response to new growth, because they prefer young leaves, but will also infest fruit. Citrus red mites feed on the cells of leaves and fruit. Damage to foliage produces a pale stippling that is visible on the upper leaf surface. Stippling of the green fruit disappears when the fruit change color. When large populations feed on fruit, the silvering may persist. They have multiple generations and are fed on by a large predator population of mites, lady beetles, lace wings, mantispids and the six-spotted thrip.

### Two-spotted Mites (*Tetranychus urticae*)

The two-spotted mite has a small, pale yellow body with dark spots on each side of its body. All stages overwinter on the trees and fruit. If winter is mild, they will feed through the season and increase normally. Heavy infestations produce a webbing that may cover several fruit and foliage. Eggs are clear and become opaque before hatching. The mites molt three times after hatching. If conditions are optimum, they can complete development in seven days.

Damage from feeding results in yellowing or stippling, producing a grayish cast on the foliage. Mites, eggs, cast skins and egg shells can be observed along the veins on the underside of the leaves.

Predators help reduce populations. Predatory mites, lady beetles, the six-spotted thrip and the minute pirate bugs are effective in keeping populations down. Heavy spraying will reduce predators and create mite problems.

### Citrus Rust Mites (*Phyllocopturta oleivora* (Ashm.))

Rust mites are very small. They are a deep yellow and wedge shaped. A generation may be completed in one to 32 weeks. Rust mites feed on the outside exposed surface of fruit 1/2 inch or larger. The feeding destroys rind cells and causes browning on mature oranges and blackening on green fruit. Most damage occurs in early spring to early summer, although damage is possible throughout the year in groves not treated for other pests. The mites have no specific predators, but other mite predators will feed on them.

In early spring, check leaves for mites and check fruit after it reaches 1/2 inch in size.

### Mealy Bugs

Mealy bugs are soft, oval, flat, distinctly segmented and covered with white or mealy wax that extends into spikes along the abdomen and posterior end. The citrus mealy bug has a yellow-orange body covered with a powdery wax. The waxy spikes are not very long on the abdomen or posterior. The Comstock mealy bug has a thicker wax covering, and the wax spikes on the abdomen and posterior are prominent.

The female lays several hundred eggs within 10 to 20 days in waxy egg sacks attached to the plant and fruit. There are two to three overlapping generations a year. They overwinter as eggs or in various stages, weather permitting.

Since they feed continuously, they excrete the excess sugary plant fluids onto the plant. This creates an ideal food for bees and wasps and an excellent medium for the growth of several species of fungi that develop into a black mat-like growth on the plants known as sooty mold.

### Armored Scales

**Florida Red Scale (*Chrysomphalus aonidum* (L.)):** The adult female scale is circular in outline, about 1/12 inch in diameter, dark reddish-brown with a conspicuous light yellow center. The female lays bright yellow eggs that hatch into lemon yellow oval-shaped crawlers. The eggs are deposited under the female's shell-like cover. There are four to five generations a year.

They feed on the exposed surfaces of the leaves and fruit. Injury appears as yellow spots on the leaves and fruit, which can be followed by heavy leaf and fruit drop in dense populations. These exposed branches can be killed by cold weather in winter and early spring. Inspections in orchards should be made from late spring to fall.

**Yellow Scale (*Aonidiella citrina* (Coq.)):** The yellow scale can be distinguished from the red scale by the light yellow color of its armor shell. Like the red scale, it has a circular shell, but is yellow to light orange and much flatter than the red scale. The scale body is visible through the armor and is yellow and kidney shaped. The female gives birth to live young.

Injury and feeding locations and preferences are similar to the red scale. It also has multiple generations each year.

**Purple Scale (*Lepidosaphes beckii* (Newm.)) and Glover Scale (*Lepidosaphes gloverii* (Pack.)):** The female purple scale lays grayish eggs in a sac-like enclosure under her shell. Glover scale eggs are pink and are found in two rows. Crawlers of both are off-white and oval with a posterior brown tip. There are usually three generations a year with peaks in March-April, June-July and September-October. They feed on the foliage, fruit and wood of the trees and are often overlooked because they live primarily on the inside of the tree. They like the shady areas such as the undersurface of leaves and collect along the mid-rib on the underside of the foliage. Yellow chlorotic areas on the foliage can cause defoliation and twig death. Injury to the fruit may cause fruit drop as well as green spots that cannot be removed.

Both females are long and tapered. The purple scale is wider, somewhat larger and darker. The Glover scale is very slender and elongated and is lighter or tan.

## Soft Scales

**Cottony Cushion Scale (*Icerya purchasi* Mask.):** This scale is easily recognized. The mature female has a bright orange-red, yellow or brown body that is practically or entirely covered with yellowish or white wax. They produce a fluted egg sac that is usually 2 to 2.5 times longer than the body. This sac may contain up to 1,000 red eggs. Depending upon the temperature, they may hatch in a few days or over two months. Newly hatched nymphs are bright red with dark antennae and thin brown legs. This is the primary dispersal stage and can be windblown, crawl or hitchhike on other animals.

The cottony cushion scale can severely damage trees and nursery stock. Decreased vitality, fruit drop and defoliation result from scale feeding. Most damage is caused by the feeding of the immature stages

on the foliage. Older nymphs migrate to the larger twigs and the adults to the larger branches and trunk. This scale has been known to inject toxins, which can cause tree death, into the tree, while feeding. Damage may also result from the excretion of honeydew and the development of sooty mold on the foliage and fruit, reducing photosynthesis.

**Florida Wax Scale (*Ceroplastes floridensis* Comst.):** These soft scales are covered with a dense layer of wax. The female beneath is bright pink. They feed on all parts of the plant and excrete large amounts of honeydew. They have three generations each year, and each female can lay 80 pink oval eggs at maturity. Upon hatching, the pink crawlers move about the plant and, once settled, begin to secrete the white wax. Young instars have a reddish body with white wax rays extending from the margins and dorsal areas of the body. The wax completely covers the body by the late third instar. On the foliage, the scales appear to collect along the main veins of the leaf and cover the leaf in dense populations. They are not known to cause any appreciable damage but, with the honeydew excretions, can cause the plant to look bad and grow erratically. Adults killed by sprays will not easily fall from the plant because the wax will hold them to the surface. Dead scales will fall off with time; heavy rainfall will assist in removal. Control is noted by reduced numbers and no movement to new growth.

## White Flies/Black Flies

The citrus white fly, the woolly white fly, the cloudy white fly and the black fly are all present in Louisiana. These are piercing, sucking feeders in both the nymph and adult stages. The eggs are laid individually by the white flies and are usually yellow when laid and turn dark before hatching. The black fly lays its eggs in a spiral fashion, making them easy to detect. The nymphs are clear when they first hatch and gradually change color as they mature, depending on species. The eggs are usually placed on the underside of the leaf surface, and the nymphs develop there. The nymphs develop through three instars and then pupate. These empty clear shells are often mistaken for developing nymphs. Peak times for white flies are in March-April, June-July and September-October. Populations are easily managed when caught early, before multiple generations can develop.

## Leafminers

These small pests affect the foliage of each flush of growth. The moths deposit their eggs on the underside of the foliage and, upon emerging, larvae tunnel into the leaf and create winding tunnels between the upper and lower surfaces as they develop and feed. At maturity, the larvae move to the edge

of the leaf causing it to curl. This protected area is where the larvae pupate. The larval stage lasts from five to 20 days; pupation takes from six to 22 days. Adult females emerge in the morning and lay their eggs at night. Total development time takes from 13 to 52 days, depending on weather and temperature. Adults are short lived. There are multiple generations which can occur every three weeks, depending on the environmental conditions.

### Leaf Miner Injury



### Giant Swallow Tail/ Orange-dog Caterpillar (*Papilio cresphontes* Cram.)

The orange-dog caterpillar is often a pest of young trees. One or two of these caterpillars can completely strip a young tree of its foliage. The larva appears as bird droppings on the foliage and stems when small. When disturbed, they evert a pair of orange glands from the base of the head. This is caused by blood pressure, and these horn-like glands have a very pungent odor. The odor is used as a defense mechanism against predators. The adult butterfly is called the giant swallowtail. It is black with a series of yellow spots that form bands in both the fore and hind wings.

### Leaf-footed Bugs

The leaf-footed bug is widespread and a pest of many crops including fruits, vegetables, grains, nuts and ornamentals. It is a major pest of citrus where feeding on ripening fruit causes premature color break and fruit drop. Adults will fly distances and enter orange groves at bloom time to feed on buds and young shoots. Later adult bugs will attack the ripening fruit, causing drop. Puncturing the fruit allows pathogens easy access and promotes rotting. Injury usually occurs as the fruit matures in the fall. The leaf-footed bugs will aggregate in large colonies on individual trees while neighboring trees are completely free.

## Aphids

Aphids or plant lice are injurious to young trees and can cause severe leaf curling to new growth on trees, particularly in the fall. Five different species of aphids are found on citrus: the green citrus aphid, the cotton aphid, the melon aphid, the black citrus aphid and the green peach aphid. All aphids feed with piercing, sucking mouthparts and excrete the excess sugars, honeydew, which promotes sooty mold development. Aphids can be recognized by the pair of little tubes located on the rear of the abdomen. These are called cornicles, and only aphids have them. Their feeding on the foliage causes the leaves to roll and curl up and become disfigured. This is usually done on the young tender growth, both in the spring and on the last flush of the trees in the fall. Aphids are heavily fed on by predators, mostly lady beetles, and by various parasites.

## pH

pH is a factor of the acidity or alkalinity of your water. It is important for your water to be on the acid side when spraying insecticides. Most insecticides are acid-forming materials. If mixed in an alkaline water solution, they can be broken down before you can spray them on your crop.

The average pH of water in the state is 8.3, while the optimum range for insecticides is between 5.5 and 6.5. It is best to check your water with a digital pH pen. pH can be adjusted by adding a buffer before adding the insecticide. Add buffer, check pH and repeat until the proper range is reached. Then add insecticide and spray.

Using the correct pH allows the insecticide to give you the proper knockdown of the pest and extended residual for proper insect management. Spraying without adjusting pH can cause you to spray more and lead to development of insect tolerance or resistance to the insecticide used.

Spraying with the proper pH will give better control of the pest with fewer sprays – thus saving you time and money, as well as being safer for the environment.

## Disease Control

Citrus trees and their fruit are subject to several diseases. Only the most common ones are described.

### Scab

Scab is primarily a disease of satsuma, grapefruit and lemons. It does not affect oranges. Scab affects fruit, leaves and young shoots, causing irregular raised corky, scabby wart-like outgrowths. Severely scabbed leaves and fruit become misshapen and distorted. The rind of scabbed fruit is thick and puffy. Scab infection occurs on young growth in the spring. Copper fungicides, tribasic copper sulfate or Kocide sprayed after bloom when the fruit is pea size will control scab.

### Scab Injury




### Melanose

Melanose is caused by a fungus that also affects leaves, shoots and fruit. It forms numerous dark brown dots or spots on the leaves, young shoots and fruit. These spots are at first sunken but later become slightly raised but not as much as scab. The spots may cover one side of the surface of the fruit, or they may run in streaks to form a tear stain-like pattern. Melanose infection occurs on young, tender growth. The fruit becomes progressively resistant with age.

Two methods are used for the control of Melanose, sanitation and spraying.

1. Sanitation. The Melanose fungus produces spores on dead twigs and branches. Pruning and burning the dead wood helps to control this disease by eliminating much of the source of infection.
2. Spraying. To control Melanose on the fruit, spray one to three weeks after bloom and fruit set when the fruit is pea sized with a copper fungicide, tribasic copper sulfate or Kocide. (See Citrus Spray Schedule.)

### Rust Mites - Melanose Injury



### Foot Rot

Foot rot is the most frequently encountered disease on the trunks of citrus trees in Louisiana. The disease is caused by a soil-borne fungus. Foot rot produces motile spores that usually invade the trunk at the bud union. Wet conditions during the spring favor fruit rot development. Initial symptoms include water soaking of the bark that appears as a dark spot on the trunk. At first the bark appears firm, but with age it becomes cracked and may shred as it dries. Gumming often accompanies advanced stages of foot rot. Maintaining adequate drainage by planting trees on raised beds and pruning lower limbs off the ground to allow air movement around the tree are the best controls for the homeowners. Use recommended herbicides to prevent weed growth.

### Sooty Mold

The sooty mold fungus is not a parasitic organism. It does not penetrate the tissue of the plant, but grows superficially on the honeydew excretions of white flies, aphids, mealy bugs and scale insects. Sooty mold can cause damage by preventing the sunlight from reaching the leaf and by making the fruit black and unattractive. Fruit covered with sooty mold is smaller and does not color well.

Control sooty mold indirectly by controlling the insect white flies and scales that excrete the honeydew on which the sooty mold fungus grows. When the insects are controlled, the sooty mold disappears.

### Sooty Mold Injury



### Lichens

Growth of different kinds of lichens often occurs on trunks, branches and sometimes on leaves of citrus trees. Lichen growth is less abundant on healthy, vigorous trees than on neglected, weakened trees that are growing poorly. The presence of lichens is often blamed for the unthrifty condition of the trees, but the reverse is true; the lichen's growth is abundant because the tree is unthrifty from some other cause. Lichens are generally considered harmless. They are not parasitic, do not invade the tissue of the bark and cause no damage to the tree. Lichens are easily controlled by spraying with copper fungicides.

### Fruit Rots

Fruit rots are usually not a problem in home orchards. However, Green Mold caused by *Penicillium* and Brown Rot caused by *Phytophthora* can cause problems in wet seasons or if the fruit remains on the tree too long. These fungi easily invade overripe or bruised or injured fruit. Fruit rots can be controlled by following a recommended spray program, using good sanitation, pruning low limbs to prevent fruit from touching the ground and drying fruit immediately after harvesting.

### Puffy Fruits

Puffy and misshapen fruit are mostly a problem on young vigorous growing satsuma trees. As the tree becomes older, the occurrence of puffy fruit decreases.

Puffy fruit on older trees are the result of fruit set on late blooms during periods of warm weather. Little can be done to prevent puffy fruit. Good growing conditions, proper fertility and pest control will help to reduce the amount of this condition.

## Freeze Protection

The major problem for home citrus production is survival of trees after hard freezes. Temperatures in the mid to low teens for 5 or more hours is necessary to kill trees. Temperatures in the low 20s for more than 5 hours will damage the fruit. The best way to reduce freeze damage is to maintain healthy trees. Weak trees that show diseases and insect damage or nutritional deficiencies are more susceptible to freeze damage than healthy trees. Cultural practices that induce and maintain dormancy in the winter will also help trees survive during freezes. They include no late summer or fall fertilization or pruning.

Maintaining bare ground free of mulch and grass under citrus trees is a practice often overlooked by homeowners. Bare ground under the trees is warmer during freezes than sod-covered or mulched ground. Grass and mulches prevent heat from entering the soil during the day, so less heat energy is stored in the soil for release during the night.

To protect a single tree, homeowners can construct a simple frame covered with clear plastic over the tree. Light bulbs placed near the trunk of the tree or an electric heater can raise the temperature in the frame a few degrees and protect the trees during the freeze. The structure needs to be opened by mid morning during bright sunny days to prevent the development of high temperatures that can damage the tree.

Wrapping the trunk of the tree to cover the bud union with insulation or Styrofoam will help prevent cold damage to the trunks. The top of the tree may be killed during a freeze; a tree can recover if its trunk is intact. Tree wraps work best on young trees. To prevent foot rot, treat the trunk of the trees with a copper fungicide before wrapping. The wrap should be removed in the spring to prevent the occurrence of foot rot.

Trees can also be protected from freeze injury by banking the trunk to cover the bud union of the tree with a mound of soil. A mound of soil 18 to 24 inches high extending 2 to 3 feet from the trunk of the tree is necessary to provide freeze protection. Banking needs to be done well ahead of the killing freeze. The soil must be removed from the tree in the spring to prevent foot rot. Banking and removing the soil from citrus trees are difficult tasks. It is often very difficult

for homeowners to bank trees successfully. As with wrapping, the trunk of the trees should be treated with a copper fungicide before banking.

## Salt Damage

In Louisiana's coastal parishes, saltwater intrusion into the groundwater has caused salinity problems for citrus trees. This is a serious problem that is difficult to correct.

Trees affected by salt have few leaves, bloom sparingly and bear small crops of small-sized fruit. The first signs of salt damage are the shedding of the mature leaves with the petiole of the leaf still attached to the branches. A large amount of the leaves will fall to the ground after a rain. As the damage progresses, the branches become defoliated and die back, and the canopy of the tree becomes very thin. Flowering and fruit yields are decreased.

The best defense against salt problems for home growers is to plant on a mound of soil 12 inches to 18 inches high. The mound helps to improve the drainage around the tree.

The use of underground drainage tubes (4-6 inches in diameter) has helped to minimize the damage from salt. The tubes are buried 2-feet to 3-feet deep and 5 feet to 10 feet to the side of the tree. These tubes remove water and salt away from the tree, which means this underground drainage system lowers the water table and reduces the salt in the soil where the tree is growing.

Gypsum also can be applied to the soil before planting or to the soil surface under bearing trees at the rate of 3 pounds to 5 pounds per 100 square feet.

Gypsum should be applied underneath trees every 2 to 3 years. The calcium in the gypsum combines with the salt and becomes soluble. The salt can then be leached throughout the soil by rainfall. This is a slow process, however. The calcium from the gypsum also helps combat the effects of the salt in the trees.

The use of low-salt fertilizers also is recommended. Low-salt fertilizers are made from materials with low-salt indexes such as potassium sulfate, diammonium phosphate and calcium nitrate. The use of calcium nitrate (low-salt index) is recommended for the May-June application of nitrogen. These low-salt fertilizers may be difficult to find for home growers, however, so they are encouraged to ask their local feed and seed stores and garden centers to stock low-salt fertilizers.

Splitting the amount of fertilizer applied to the trees into three applications – coming four weeks apart in February, March and April – instead of just one application in February also is recommended. These frequent applications provide the necessary nutrients with a minimum amount of salt added to the soil.

Control-released, low-salt fertilizers also can be used in February on citrus trees having problems with salt. These fertilizers release a small amount of nutrients over a period of several months. Most controlled-released fertilizers are liquid encased in a plastic coating. The coating is washed away, which then releases the liquid. The coating on the particles washes away at different rates, providing a continuous release of small amounts of fertilizer.

Keeping the soil moist under the trees by frequent light watering of the trees also will help to reduce the effects of salt on citrus trees.



*Salt Damage to Citrus Tree*

### Simplified Citrus Spray Schedule for Home Gardens

These spray applications should control most insects and diseases.

Time	Pesticide and Formulation	Amount to use for 50 gals.	1 gal.	Pest controlled	Limitations Good coverage needed for control
Post-bloom spray (when 75% of petals have fallen)	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	scale, whitefly, mealy bug, spider mites, rust mites	0 days to harvest
	OR				
	Malathion (57% EG)	1 pt.	2 tsp.	scale, whitefly, mealy bug	Do not apply Malathion to plants in full bloom. 7 days to harvest
	+				
	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	spider mites, rust mites	
	OR				
Post Bloom	Kelthane MF (4 EC)	1/2 pt.	1 tsp.	spider mites, rust mites	7 days to harvest
	+				
	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	scale, whitefly, mealy bugs, spider mites, rust mites	
	OR				
	Dimethoate 2.67 EC	3/4 - 1 pt.	2 tsp.	scale, whitefly	Do not apply during bloom. Make no more than 2 applications to mature fruit. 15 days to harvest.
	+				
Post Bloom	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	scale, whitefly, mealy bugs, spider mites	
	Vendex 50 WP	2-4 oz.	1 tsp.	spider mites, rust mites	4 applications per season. Do not use on tangerines, tangelos or Reed, Webb or Red Blush grapefruit. 7 days until harvest.
	Spinosid	4-6 ozs.	1 tsp.	leafminers	7 days to harvest. Applications should be made at each flush of growth.
First application of fixed copper when fruit is pea size	Fixed Copper (Liquid copper fungicides, Tribasic Copper Sulfate, Kocide DF, Kocide 2000	Mfg. Label	Mfg. Label	scab, melanose	Spray fixed copper at pea-sized fruit and spray again in 2 weeks. Do not mix copper with other pesticides. Do not use Bluestone on citrus. Scab - satsuma Melanose - oranges
Second application of fixed copper, 2 weeks after first application	Fixer Copper (Liquid copper fungicides, Tribasic Copper Sulfate, Kocide DF, Kocide 2000	Mfg. Label	Mfg. Label	scab, melanose	Make second application of fixed copper 2 weeks after first application. Do not mix copper with other pesticides. Do not use Bluestone on citrus. Scab - satsuma Melanose - oranges

Time	Pesticide and Formulation	Amount to use for		Pest controlled	Limitations Good coverage needed for control	
		50 gals.	1 gal.			
Summer spray May-August	Spinosid	4 - 6 oz.	1 tsp.	leafminers, caterpillars	7 days before harvest	
	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	scales, mealy bugs, whitefly, spider mites, rust mites	0 day to harvest	
	OR					
	Summer Oil Emulsion	1/2 gal.	5 T.	scales, mealy bugs, whitefly, spider mites, rust mites	0 day to harvest	
	OR					
	Malathion 57% EC + Sun Spray Ultra Fine Oil	1 pt.	2 tsp.	scales, mealy bugs, whitefly, spider mites, rust mites	7 days until harvest	
	OR					
Dimethoate 267 EC + Sun Spray Ultra Fine Oil	3/4 - 1 pt.	2 tsp.	scales, mealy bugs, whitefly, spider mites, rust mites	15 days to harvest. Make no more than 2 applications to mature fruit.		
OR						
OR	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	spider mites, rust mites		
Fall Sept. - Nov.	Spinosid	4 - 6 oz.	1 tsp.	leafminers, caterpillars	7 days before harvest	
	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	3-4 T.	scales, mealy bugs, whitefly, spider mites, rust mites	0 day to harvest	
	OR					
	Malathion 57% EC + Sun Spray Ultra Fine Oil	1 pt.	2 tsp.	scales, mealy bugs, whitefly, spider mites, rust mites	7 days to harvest	
	OR					
	Dimethoate 2.76 EC + Sun Spray Ultra Fine Oil	3/4 - 1 pt.	2 tsp.	scales, mealy bugs, whitefly, spider mites, rust mites	15 days to harvest, make no more than 2 applications to mature fruit.	
	OR					
	OR	Sun Spray Ultra Fine Oil	1/2 - 1 gal.	2-4 T.	spider mites, rust mites	
	OR					
	Kelthane MF (4EC) + Sun Spray Ultra Fine Oil	1/2 pt.	1 tsp.	spider mites, rust mites, scale, mealy bugs, whitefly	7 days to harvest	
OR						
OR	Vendex 50 WP	2-4 oz.	1 tsp.	spider mites	7 days to harvest, 4 applications per season. Do not use on tangerines, tangelos or Reed, Webb or Red Blush grapefruit.	

### **Acknowledgments**

The authors wish to acknowledge the following individuals for their contributions to this publication:

Wayne Bourgeois, Professor, Citrus Research Station

Alan Vaughn, County Agent, Plaquemines Parish

Stuart Gauthier, County Agent, Vermilion Parish

Bobby Fletcher Jr., Area Agent, Terrebonne/Lafourche Parishes



**Visit our Web site: [www.lsuagcenter.com](http://www.lsuagcenter.com)**

### **Authors**

James Boudreaux, Professor (Horticulture)

Don Ferrin, Assistant Professor (Plant Pathology)

Dale Pollet, Professor (Entomology)

Ken Whitam, Professor (Plant Pathology - retired)

### **Louisiana State University Agricultural Center**

William B. Richardson, Chancellor

### **Louisiana Agricultural Experiment Station**

David J. Boethel, Vice Chancellor and Director

### **Louisiana Cooperative Extension Service**

Paul D. Coreil, Vice Chancellor and Director

**Pub. 1234**

**(7M)**

**02/07 Rev.**

Issued in furtherance of Cooperative Extension work, Act. of Congress of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. The Louisiana Cooperative Extension Service provides equal opportunities in programs and employment.