

LOUISIANA RECOMMENDATIONS FOR CONTROL OF INSECTS ON PASTURES AND FORAGE CROPS

Insect	Crop	Insecticide	lbs. Active Ingredient/Acre	Limitations
Spider mites	Clovers	Malathion*	1 lb	Most insecticides are toxic to bees. Avoid treating pollinating forages when bees are active.
Cutworms	Clovers	Sevin/carbaryl*	1.0-1.5 lbs	<u>Pre-harvest treatment interval for PASTURE GRASS, CLOVER, and VETCHES.</u>
Armyworms and Fall armyworms	Pasture grasses and hay crops	Sevin/carbaryl Lannate Methyl parathion Tracer Treat for 1 or more worms per sweep or 1-2 or more per square foot.	0.5 lb 0.225 lb 0.5 lb 0.031- 0.062 lb.	<u>Lannate:</u> 7-day PHI-grazing or harvest. REI: 48 hrs. <u>Malathion:</u> 0-day PHI-grazing or harvest. <u>Methyl parathion/Pennacap M:</u> 15-day PHI-grazing or harvest. REI: 4 days
Chinch bugs	Pasture grasses Forage Sorghums	Carbaryl (Sevin) Carbaryl (Sevin)	1.5 lbs 1.0-2.0 lbs	<u>Sevin: (Pastures)</u> 14-day PHI-grazing or harvest for improved pasture; 0 days for rangeland (no improvements). Do not apply more than 1.5 lbs. AI/acre/season. REI: 12 hrs.
Clover head weevils	Clovers	Methyl parathion Examine seed heads. Treat when 10%-20% of seed heads are infested with larvae.	0.5-1.0 lb	<u>Tracer:</u> Toxic to bees and mollusks. Pre-harvest interval: 3 days for hay/fodder; 0 for fodder. REI: 4 hrs.
Spittlebugs	Bermuda grass	No chemical control is recommended. Prevent a dense mat of grass from forming by grazing or by close mowing and raking.		<u>Pre-harvest treatment interval for ALFALFA.</u> <u>Dimethoate:</u> 10-day PHI-grazing or harvest and only one application per cutting. REI: 48 hrs.
Aphids	Clovers Vetches Grasses	Malathion Methyl parathion	1 lb 0.25 lb	<u>Furadan:</u> 7-day PHI for 0.25 lb. AI/acre rate, 14 days for 0.5 lb AI/acre rate. Do not apply more than once per cutting. REI: 14 days.
Plant bugs	Clovers and Legumes (seed crop only)	Methyl parathion	0.25-0.5 lb	<u>Imidan:</u> 7-day PHI-grazing or harvest. REI: 24 hrs.
Leaf hoppers	Pasture grasses and Alfalfa	Carbaryl (Sevin) Close graze or harvest Bermuda grass pastures at 6-inch height to reduce leafhopper/plant hopper populations	1 lb	<u>Methyl parathion/Pennacap M:</u> 15-day PHI -grazing or harvest. REI: 4 days. <u>Sevin:</u> 7-day PHI-grazing or harvest and only one application per cutting. Do not apply more than 1.5 lbs. AI/acre/cutting. REI: 12 hrs.

**LOUISIANA RECOMMENDATIONS FOR CONTROL OF
INSECTS ON PASTURES AND FORAGE CROPS (cont'd)**

Insect	Crop	Insecticide	lbs. Active Ingredient/Acre	Limitations
				Lorsban: 14 days for 0.5 lb AI/acre rate, grazing or cutting, and only one application per cutting. REI: 24 hrs.
Spotted alfalfa aphids	Alfalfa	Malathion	0.75 lb	Baythroid: Extremely toxic to fish and aquatic invertebrates. Avoid drift, runoff, and direct application to water; 7-day pre-harvest and pre-grazing interval. REI: 12 hrs.
		Methyl parathion	0.25 lb	
		Pennacap-M	0.25 lb	
		Dimethoate	0.25-0.5 lb	
		Mustang Max	0.014-0.025	
Alfalfa weevils	Alfalfa (Treat when 50% of growing tips have feeding sign)	Methyl parathion	0.5 lb	Mustang Max: Pre-harvest interval, 3 days for cutting or grazing, 7 days for seed. Do not apply more than 0.025 lb AI per cutting or 0.075 lb AI per season. REI: 24 hrs.
		Furadan 4F	0.25-0.5 lb	
		Imidan	0.9 lb	
		Lorsban	0.5 lb	
		Baythroid	0.025-0.044	
		Pennacap-M	0.5 lb	
		Mustang Max	0.014-0.025	
Warrior T	0.02-0.03			

Imported fire ant Apply fire ant baits when soil temperature is 60°F or higher, and rain is not forecast.	Pasture grasses	Amdro Pro	1.0-1.5 lbs bait/acre broadcast or 5 TBS/mound	Karate Z: Pre-harvest interval 7 days for hay and 1 day for forage. Do not apply more than 0.03 lb. AI/acre/cutting or more 0.12 lb. AI/acre/season. REI: 24 hrs.
		Carbaryl (Sevin)	Use as a mound drench as per label instructions.	
		Extinguish (methoprene)	1.0-1.5 lbs bait/acre broadcast or 3-5 TBS/mound	
		Logic* (fenoxycarb) (Horse pasture only)	1.0-1.5 lbs/acre or 1-3 TBS/mound	
		Justice (spinosad)	4-6 TBS/mound Do not broadcast.	
		Esteem (pyriproxifen)	1.5-2.0 lbs./acre	
				Amdro: 7-day pre-harvest interval for cutting and baling hay. Maximum applications/year: 4
				Extinguish: 0 days.
				Esteem: Pre-harvest interval: 24 hrs.
				Logic: for use on HORSE PASTURES only.
				Justice: No grazing restrictions.

A reduction of approximately 50% in mound number and mound size may be obtained by dragging pastures and disrupting ant mounds 2 times per winter shortly before and/or during the time the air temperature is 32°F or lower.

* Horse pastures only.